

CLAIMS

5 v b
a 2
1. A pharmaceutical composition or a dietary supplement comprising:

- i) an extract or concentrate of *Butyrospermum parkii* containing at least 5% (w/w) of a
5 *Butyrospermum*-triterpene fraction comprising:
- at least 2% (w/w) lupeol;
- at least 2% (w/w) α -amyirin and/or β -amyirin;
- at least 2% (w/w) butyrospermol; and
- optionally at least 1% germanicol, dammaradienol, 24-methylene-dammarenol and/or
10 parkeol,

wherein said triterpenes may be in the form of free alcohols or esters thereof, especially cinnamic acid, acetic acid or fatty acid esters; and

- 15 ii) optionally a pharmaceutically acceptable carrier.

2. A pharmaceutical composition or a dietary supplement comprising:

- i) an extract or concentrate of *Butyrospermum parkii* containing at least 5% (w/w) of a
20 *Butyrospermum*-triterpene fraction comprising:
- 10-40% (w/w) lupeol;
- 10-40% (w/w) α -amyirin and/or β -amyirin;
- 10-40% (w/w) butyrospermol; and
- optionally 2-30% germanicol, dammaradienol, 24-methylene-dammarenol and/or
25 parkeol,

wherein said triterpenes may be in the form of free alcohols or esters thereof, especially cinnamic acid, acetic acid or fatty acid esters; and

- 30 ii) optionally a pharmaceutically acceptable carrier.

3. A pharmaceutical composition or dietary supplement according to claim 1 or 2, where the extract or concentrate of *Butyrospermum parkii* further comprises a sterol fraction comprising at least one sterol selected from the group consisting of stigmasterol, av-

nasterol, 24-methyl-cholest-7-enol, karitesterol A, karitesterol B and α -spinasterol, wherein said sterols may be in the form of free alcohols or esters thereof, especially cinnamic acid, acetic acid or fatty acid esters.

- cont*
cl
- B*
B
- 5 4. A pharmaceutical composition or dietary supplement according to ~~any of the preceding claims~~ *claim 1*, wherein the Butyrospermum-triterpene fraction optionally together with the sterol fraction comprises up to 100% (w/w) of the extract or concentrate of Butyrospermum parkii.
- 10 5. A pharmaceutical composition or dietary supplement according to any of claims 3 or 4, wherein the ratio between the Butyrospermum-triterpene fraction and the sterol fraction is in the range of 1:100 to 500:1 (w/w).
- B*
B
6. A pharmaceutical composition or dietary supplement according to ~~any of the preceding claims~~ *claim 1*, which further comprises an extract of Calendula officinalis.
- 15 7. A pharmaceutical composition according to ~~any of the preceding claims~~ *claim 1* for systemic administration.
- B*
B
- 20 8. A pharmaceutical composition according to ~~any of claims 1 to 6~~ *claim 1* for topical administration, wherein the pharmaceutical composition contains at least 5% (w/w) of the Butyrospermum-triterpene fraction.
9. A pharmaceutical composition according to claim 8, wherein the pharmaceutical
- 25 composition is formulated as a fluid, ointment, gel, liniment, emulsion (e.g. cream or lotion) or spray (e.g. aerosol).
- B*
10. The use of a composition according to ~~any of claims 1 to 9~~ *claim 1* for the preparation of a medicament or a dietary supplement for immunomodulation in a mammal.
- 30
11. The use of a composition according to ~~any of claims 1 to 9~~ *claim 1* for the preparation of a medicament or a dietary supplement for the suppression of hypersensitivity and/or inflammatory reaction in a mammal.
- B*

12. The use of a composition according to claim 11 for the preparation of a medicament for the treatment or prevention of inflammation or hypersensitivity of the skin or mucous membranes in a mammal.

- 5 13. The use according to claim 11 or 12 for the preparation of a medicament or a dietary supplement for the treatment or prevention of autoimmune disease and/or chronic inflammatory disease in a mammal.

14. The use according to claim 13 for the preparation of a medicament or a dietary
10 supplement for the treatment or prevention of psoriasis, atopic dermatitis, contact dermatitis, Crohn's disease, ulcerative colitis, rheumatoid arthritis or osteoarthritis in a mammal.

15. The use of a composition according to ~~any of claims 1 to 9~~ ^{claim 1} for the preparation of a me-
15 dicament or a dietary supplement for the alleviation of pain in a mammal.

16. The use of a composition according to ~~any of claims 1 to 9~~ ^{claim 1} for the preparation of a me-
dicament or a dietary supplement for the treatment or prevention of prostatitis and/or be-
nign prostatic hypertrophy.

- 20 17. A method for the treatment or prevention of hypersensitivity or inflammation in a mam-
mal, characterised by administering a composition according to ~~any of claims 1 to 9~~ ^{claim 1} to
said mammal.

- 25 18. A method for the treatment or prevention of inflammation or hypersensitivity of the skin
or mucous membranes of a mammal, characterised by administering a composition ac-
cording to ~~any of claims 1 to 9~~ ^{claim 1} to said mammal.

19. A method for the treatment or prevention of an autoimmune disorder and/or a chronic
30 inflammatory disorder in a mammal, characterised by administering a mixture according to
~~any of claims 1 to 9~~ ^{claim 1} to said mammal.

20. A method for the treatment or prevention of psoriasis, atopic eczema, contact
dermatitis, Crohn's disease, ulcerative colitis, rheumatoid arthritis and/or osteoarthritis in a

mammal, characterised by administering a mixture according to ~~any of claims 1 to 9~~ ^{claim 1} to said mammal.

21. A method for the treatment or prevention of pain in a mammal, characterised by administering a mixture according to ~~any of claims 1 to 9~~ ^{claim 1} to said mammal.

22. A method for the treatment or prevention of prostatitis or benign prostatic hypertrophy in a mammal, characterised by administering a mixture according to ~~any of claims 1 to 9~~ ^{claim 1} to said mammal.

23. A method for the preparation of a composition according to ~~any of claims 1 to 9~~ ^{claim 1}, characterised by obtaining an extract or a concentrate of *Butyrospermum parkii*, said extract or concentrate containing at least 5% (w/w) of a Butyrospermum-triterpene fraction comprising:

- at least 2% (w/w) lupeol;
- at least 2% (w/w) α -amyirin and/or β -amyirin;
- at least 2% (w/w) butyrospermol; and
- optionally at least 1% germanicol, dammaradienol, 24-methylene-dammarenol and/or parkeol,

wherein said triterpenes may be in the form of free alcohols or esters thereof, especially cinnamic acid, acetic acid or fatty acid esters; and

24. A method according to claim 22, wherein the extract or concentrate of *Butyrospermum parkii* further comprises a sterol fraction comprising at least one sterol selected from the group consisting of stigmasterol, avanasterol, 24-methyl-cholest-7-enol, karitesterol A, karitesterol B and α -spinasterol, wherein said sterols may be in the form of free alcohols or esters thereof, especially cinnamic acid, acetic acid or fatty acid esters.

25. A method according to claim 22 or 23, wherein said extract or concentrate of *Butyrospermum parkii* is further mixed with a pharmaceutically acceptable carrier.